

# DIRECTIONS FOR USE

# PEROXIGARD™

ACCELERATED HYDROGEN PEROXIDE™

Product	Site Area	Dilution	Directions For Use	Protocol
<b>Ready-To-Use Surface Cleaner and Disinfectant (1 L)</b>  For use as a disinfectant	<b>Environmental Surfaces and Equipment</b>  Walls, counter tops, hatchery tanks, graders, vaccination equipment, nets and foot baths.	No dilution required	<b>Disinfection</b> <b>Surfaces:</b> Apply to surface <ul style="list-style-type: none"> <li>• Let sit wet for <b>5 min</b></li> <li>• Wipe once using one side of cloth then turn cloth over and wipe</li> <li>• Wipe dry or rinse</li> </ul> <b>Nets:</b> Change frequently <b>Foot baths:</b> Change frequently or when heavily soiled	Heavily soiled surfaces require cleaning prior to disinfection
<b>Ready-To-Use Surface Cleaner and Disinfectant (1 L)</b>  For use as a cleaner and sanitizer	<b>Environmental Surfaces and Equipment</b>  Walls, counter tops, hatchery tanks, graders, vaccination equipment and vehicles.	No dilution required	<b>Cleaning and Sanitizing</b> <ul style="list-style-type: none"> <li>• Apply to surface</li> <li>• Let sit wet for <b>30 sec</b></li> <li>• Wipe once using one side of cloth then turn cloth over and wipe</li> <li>• Wipe dry</li> <li>• No rinsing required</li> </ul>	Remove organic debris prior to cleaning process.  Heavily soiled surfaces should be cleaned twice. Food contact surfaces require rinsing with potable water.
<b>Concentrated Surface Cleaner and Disinfectant (4 L, 20 L, 200 L)</b>  For use as a disinfectant	<b>Environmental Surfaces and Equipment</b>  Walls, counter tops, hatchery tanks, graders, vaccination equipment, nets and foot baths.	<b>1:16 dilution</b>  Add 2 pumps (2 oz/ 64 ml) of concentrate to an empty Ready-To-Use spray bottle and fill with water (32 oz/1 L)	<b>Disinfection</b> <b>Surfaces:</b> Apply to surface <ul style="list-style-type: none"> <li>• Let sit wet for <b>5 min</b></li> <li>• Wipe once using one side of cloth then turn cloth over and wipe</li> <li>• Wipe dry or rinse</li> </ul> <b>Nets:</b> Change frequently <b>Foot baths:</b> Change frequently or when heavily soiled	Heavily soiled surfaces require cleaning prior to disinfection
<b>Concentrated Surface Cleaner and Disinfectant (4 L, 20 L, 200 L)</b>  For use as a cleaner and sanitizer	<b>Environmental Surfaces and Equipment</b>  Walls, counter tops, hatchery tanks, graders, vaccination equipment and vehicles.	<b>1:16 dilution</b>  Add 2 pumps (2 oz/ 64 ml) of concentrate to an empty Ready-To-Use spray bottle and fill with water (32 oz/1 L)	<b>Cleaning and Sanitizing</b> <ul style="list-style-type: none"> <li>• Apply to surface</li> <li>• Let sit wet for <b>30 sec</b></li> <li>• Wipe once using one side of cloth then turn cloth over and wipe</li> <li>• Wipe dry</li> <li>• No rinsing required</li> </ul>	Remove organic debris prior to cleaning process.  Heavily soiled surfaces should be cleaned twice. Food contact surfaces require rinsing with potable water.
<b>Concentrated Surface Cleaner and Disinfectant (4 L, 20 L, 200 L)</b>  For cleaning floors and walls	<b>For Cleaning Floors and Walls</b>  Egg incubation rooms, quarantine facilities, general hatchery and farm use.	<b>1:64 dilution</b>  Add 2½ pumps (2.5 oz/80 ml) of concentrate per gallon/4.5 L of water	<b>Cleaning of Floors and Walls</b> <ul style="list-style-type: none"> <li>• Wash surfaces</li> <li>• Let sit wet for 5 min</li> </ul>	Remove organic debris prior to cleaning process.

Accelerated Hydrogen Peroxide is a registered trademark and patented technology of Virox Technologies Inc. used under license by Bayer Inc. Peroxigard is a trademark of Bayer Inc. Bayer and the Bayer Cross are registered trademarks of Bayer AG, used under license by Bayer Inc.



+1 250 652-4482  
 Toll free (Canada) 1-800-667-5062  
 e-mail peroxigard@bayotek.com  
 www.bayotek.com

Virucidal

Bactericidal

Broad Spectrum Sanitizer